

CHEMISTRY

Data Booklet *Updated 2010*





Government
of Alberta ■

Alberta ■
Freedom To Create. Spirit To Achieve.



10	11	12	13	14	15	16	17	18
----	----	----	----	----	----	----	----	----

Legend for Elements

	Metallic solids		Gases
	Non-metallic solids		Liquids

Note: The legend denotes the physical state of the elements at exactly 101.325 kPa and 298.15 K.

Key

Atomic number	26	55.85	Atomic molar mass (g/mol)*
		3+, 2+	Most stable ion charges
Electronegativity	1.8		
Symbol	Fe		
Name	iron		

* Based on $^{12}_6\text{C}$
 () Indicates mass of the most stable isotope

							2	4.00									
							He	helium									
		5	10.81	6	12.01	7	14.01	8	16.00	9	19.00	10	20.18				
		2.0		2.6		3.0		3.4		4.0		—					
		B	boron	C	carbon	N	nitrogen	O	oxygen	F	fluorine	Ne	neon				
		13	26.98	14	28.09	15	30.97	16	32.07	17	35.45	18	39.95				
		1.6		1.9		2.2		2.6		3.2		—					
		Al	aluminium	Si	silicon	P	phosphorus	S	sulfur	Cl	chlorine	Ar	argon				
28	58.69	29	63.55	30	65.41	31	69.72	32	72.64	33	74.92	34	78.96	35	79.90	36	83.80
1.9		1.9		1.7		1.8		2.0		2.2		2.6		3.0		—	
Ni	nickel	Cu	copper	Zn	zinc	Ga	gallium	Ge	germanium	As	arsenic	Se	selenium	Br	bromine	Kr	krypton
46	106.42	47	107.87	48	112.41	49	114.82	50	118.71	51	121.76	52	127.60	53	126.90	54	131.29
2.2		1.9		1.7		1.8		2.0		2.1		2.1		2.7		2.6	
Pd	palladium	Ag	silver	Cd	cadmium	In	indium	Sn	tin	Sb	antimony	Te	tellurium	I	iodine	Xe	xenon
78	195.08	79	196.97	80	200.59	81	204.38	82	207.2*	83	208.98	84	(209)	85	(210)	86	(222)
2.2		2.4		1.9		1.8		1.8		1.9		2.0		2.2		—	
Pt	platinum	Au	gold	Hg	mercury	Tl	thallium	Pb	lead	Bi	bismuth	Po	polonium	At	astatine	Rn	radon
110	(271)	111	(272)														
Ds	darmstadtium	Rg	roentgenium														

* The isotopic mix of naturally occurring lead is more variable than other elements, preventing precision to greater than tenths of a gram per mole.

63	151.96	64	157.25	65	158.93	66	162.50	67	164.93	68	167.26	69	168.93	70	173.04	71	174.97
—		1.2		—		1.2		1.2		1.2		1.3		—		1.0	
Eu	europium	Gd	gadolinium	Tb	terbium	Dy	dysprosium	Ho	holmium	Er	erbium	Tm	thulium	Yb	ytterbium	Lu	lutetium
95	(243)	96	(247)	97	(247)	98	(251)	99	(252)	100	(257)	101	(258)	102	(259)	103	(262)
—		—		—		—		—		—		—		—		—	
Am	americium	Cm	curium	Bk	berkelium	Cf	californium	Es	einsteinium	Fm	fermium	Md	mendelevium	No	nobelium	Lr	lawrencium

Chemistry Notation

Symbol	Term	Unit(s)
c	specific heat capacity	J/(g·°C) or J/(g·K)
E°	standard electrical potential	V or J/C
E_k	kinetic energy	kJ
E_p	potential energy	kJ
ΔH	enthalpy (heat)	kJ
$\Delta_f H^\circ$	standard molar enthalpy of formation	kJ/mol
I	current	A or C/s
K_c	equilibrium constant	—
K_a	acid ionization (dissociation) constant	—
K_b	base ionization (dissociation) constant	—
M	molar mass	g/mol
m	mass	g
n	amount of substance	mol
P	pressure	kPa
Q	charge	C
T	temperature (absolute)	K
t	temperature (Celsius)	°C
t	time	s
V	volume	L
c	amount concentration	mol/L

Symbol	Term
Δ	delta (change in)
$^\circ$	standard
[]	amount concentration

Miscellaneous

25.00 °C is equivalent to 298.15 K

Specific Heat Capacities at 298.15 K and 100.000 kPa

$$c_{\text{air}} = 1.01 \text{ J/(g}\cdot\text{°C)}$$

$$c_{\text{polystyrene foam cup}} = 1.01 \text{ J/(g}\cdot\text{°C)}$$

$$c_{\text{copper}} = 0.385 \text{ J/(g}\cdot\text{°C)}$$

$$c_{\text{aluminium}} = 0.897 \text{ J/(g}\cdot\text{°C)}$$

$$c_{\text{iron}} = 0.449 \text{ J/(g}\cdot\text{°C)}$$

$$c_{\text{tin}} = 0.227 \text{ J/(g}\cdot\text{°C)}$$

$$c_{\text{water}} = 4.19 \text{ J/(g}\cdot\text{°C)}$$

Water Autoionization Constant (Dissociation Constant)

$K_w = 1.0 \times 10^{-14}$ at 298.15 K (for ion concentrations in mol/L)

Faraday Constant

$$F = 9.65 \times 10^4 \text{ C/mol e}^-$$

Quadratic Formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Selected SI Prefixes

Prefix	Exponential Symbol	Value
tera	T	10^{12}
giga	G	10^9
mega	M	10^6
kilo	k	10^3
milli	m	10^{-3}
micro	μ	10^{-6}
nano	n	10^{-9}
pico	p	10^{-12}

Standard Molar Enthalpies of Formation at 298.15 K

Name	Formula	$\Delta_f H^\circ$ (kJ/mol)
aluminium oxide	Al ₂ O ₃ (s)	-1 675.7
ammonia	NH ₃ (g)	-45.9
ammonium chloride	NH ₄ Cl(s)	-314.4
ammonium nitrate	NH ₄ NO ₃ (s)	-365.6
barium carbonate	BaCO ₃ (s)	-1 213.0
barium chloride	BaCl ₂ (s)	-855.0
barium hydroxide	Ba(OH) ₂ (s)	-944.7
barium oxide	BaO(s)	-548.0
barium sulfate	BaSO ₄ (s)	-1 473.2
benzene	C ₆ H ₆ (l)	+49.1
butane	C ₄ H ₁₀ (g)	-125.7
calcium carbonate	CaCO ₃ (s)	-1 207.6
calcium chloride	CaCl ₂ (s)	-795.4
calcium hydroxide	Ca(OH) ₂ (s)	-985.2
calcium oxide	CaO(s)	-634.9
calcium sulfate	CaSO ₄ (s)	-1 434.5
carbon dioxide	CO ₂ (g)	-393.5
carbon monoxide	CO(g)	-110.5
chromium(III) oxide	Cr ₂ O ₃ (s)	-1 139.7
copper(I) oxide	Cu ₂ O(s)	-168.6
copper(II) oxide	CuO(s)	-157.3
copper(II) sulfate	CuSO ₄ (s)	-771.4
copper(I) sulfide	Cu ₂ S(s)	-79.5
copper(II) sulfide	CuS(s)	-53.1
dinitrogen tetroxide	N ₂ O ₄ (g)	+11.1
ethane	C ₂ H ₆ (g)	-84.0
ethanoic acid (acetic acid)	CH ₃ COOH(l)	-484.3
ethanol	C ₂ H ₅ OH(l)	-277.6
ethene (ethylene)	C ₂ H ₄ (g)	+52.4
ethyne (acetylene)	C ₂ H ₂ (g)	+227.4
glucose	C ₆ H ₁₂ O ₆ (s)	-1 273.3
hydrogen bromide	HBr(g)	-36.3
hydrogen chloride	HCl(g)	-92.3
hydrogen fluoride	HF(g)	-273.3
hydrogen iodide	HI(g)	+26.5
hydrogen perchlorate	HClO ₄ (l)	-40.6
hydrogen peroxide	H ₂ O ₂ (l)	-187.8
hydrogen sulfide	H ₂ S(g)	-20.6
iron(II) oxide	FeO(s)	-272.0
iron(III) oxide	Fe ₂ O ₃ (s)	-824.2
iron(II,III) oxide (magnetite)	Fe ₃ O ₄ (s)	-1 118.4
lead(II) bromide	PbBr ₂ (s)	-278.7
lead(II) chloride	PbCl ₂ (s)	-359.4
lead(II) oxide (red)	PbO(s)	-219.0
lead(IV) oxide	PbO ₂ (s)	-277.4
magnesium carbonate	MgCO ₃ (s)	-1 095.8
magnesium chloride	MgCl ₂ (s)	-641.3

Standard Molar Enthalpies of Formation at 298.15 K cont'd

Name	Formula	$\Delta_f H^\circ$ (kJ/mol)
magnesium hydroxide	Mg(OH) ₂ (s)	– 924.5
magnesium oxide	MgO(s)	– 601.6
magnesium sulfate	MgSO ₄ (s)	– 1 284.9
manganese(II) oxide	MnO(s)	– 385.2
manganese(IV) oxide	MnO ₂ (s)	– 520.0
mercury(II) oxide (red)	HgO(s)	– 90.8
mercury(II) sulfide (red)	HgS(s)	– 58.2
methanal (formaldehyde)	CH ₂ O(g)	– 108.6
methane	CH ₄ (g)	– 74.6
methanoic acid (formic acid)	HCOOH(l)	– 425.0
methanol	CH ₃ OH(l)	– 239.2
nickel(II) oxide	NiO(s)	– 240.6
nitric acid	HNO ₃ (l)	– 174.1
nitrogen dioxide	NO ₂ (g)	+ 33.2
nitrogen monoxide	NO(g)	+ 91.3
octane	C ₈ H ₁₈ (l)	– 250.1
pentane	C ₅ H ₁₂ (l)	– 173.5
phosphorus pentachloride	PCl ₅ (s)	– 443.5
phosphorus trichloride (liquid)	PCl ₃ (l)	– 319.7
phosphorus trichloride (vapour)	PCl ₃ (g)	– 287.0
potassium bromide	KBr(s)	– 393.8
potassium chlorate	KClO ₃ (s)	– 397.7
potassium chloride	KCl(s)	– 436.5
potassium hydroxide	KOH(s)	– 424.6
propane	C ₃ H ₈ (g)	– 103.8
silicon dioxide (α -quartz)	SiO ₂ (s)	– 910.7
silver bromide	AgBr(s)	– 100.4
silver chloride	AgCl(s)	– 127.0
silver iodide	AgI(s)	– 61.8
sodium bromide	NaBr(s)	– 361.1
sodium chloride	NaCl(s)	– 411.2
sodium hydroxide	NaOH(s)	– 425.8
sodium iodide	NaI(s)	– 287.8
sucrose	C ₁₂ H ₂₂ O ₁₁ (s)	– 2 226.1
sulfur dioxide	SO ₂ (g)	– 296.8
sulfuric acid	H ₂ SO ₄ (l)	– 814.0
sulfur trioxide (liquid)	SO ₃ (l)	– 441.0
sulfur trioxide (vapour)	SO ₃ (g)	– 395.7
tin(II) chloride	SnCl ₂ (s)	– 325.1
tin(IV) chloride	SnCl ₄ (l)	– 511.3
tin(II) oxide	SnO(s)	– 280.7
tin(IV) oxide	SnO ₂ (s)	– 577.6
water (liquid)	H ₂ O(l)	– 285.8
water (vapour)	H ₂ O(g)	– 241.8
zinc oxide	ZnO(s)	– 350.5
zinc sulfide (sphalerite)	ZnS(s)	– 206.0

Solubility of Some Common Ionic Compounds in Water at 298.15 K

Ion	Group 1 ions NH ₄ ⁺ NO ₃ ⁻ ClO ₃ ⁻ ClO ₄ ⁻ CH ₃ COO ⁻	F ⁻	Cl ⁻ Br ⁻ I ⁻	SO ₄ ²⁻	CO ₃ ²⁻ PO ₄ ³⁻ SO ₃ ²⁻	IO ₃ ⁻ OOC ⁻ COO ²⁻	OH ⁻
Solubility greater than or equal to 0.1 mol/L (very soluble)	most	most	most	most	Group 1 ions NH ₄ ⁺	Group 1 ions NH ₄ ⁺ Co(IO ₃) ₂ Fe ₂ (OOC ⁻ COO) ₃	Group 1 ions NH ₄ ⁺
Solubility less than 0.1 mol/L (slightly soluble)	RbClO ₄ CsClO ₄ AgCH ₃ COO Hg ₂ (CH ₃ COO) ₂	Li ⁺ Mg ²⁺ Ca ²⁺ Sr ²⁺ Ba ²⁺ Fe ²⁺ Hg ₂ ²⁺ Pb ²⁺	Cu ⁺ Ag ⁺ Hg ₂ ²⁺ Pb ²⁺ Tl ⁺	Ca ²⁺ Sr ²⁺ Ba ²⁺ Ag ⁺ Hg ₂ ²⁺ Pb ²⁺ Ra ²⁺	most	most	most

Note: This solubility table is only a guideline that is established using the K_{sp} values. A concentration of 0.1 mol/L corresponds to approximately 10 g/L to 30 g/L depending on molar mass. Hg₂²⁺ is a polyatomic ion of mercury.

Flame Colour of Elements

Element	Symbol	Colour
lithium	Li	red
sodium	Na	yellow
potassium	K	violet
rubidium	Rb	violet
cesium	Cs	violet
calcium	Ca	yellowish red
strontium	Sr	scarlet red
barium	Ba	yellowish green
copper	Cu	blue to green
boron	B	yellowish green
lead	Pb	blue-white

Note: The flame test can be used to determine the identity of a metal or a metal ion. Blue to green indicates a range of colours that might appear.

Table of Selected Standard Electrode Potentials*

Reduction Half-Reaction	Electrical Potential E° (V)
$F_2(g) + 2e^- \rightleftharpoons 2F^-(aq)$	+2.87
$PbO_2(s) + SO_4^{2-}(aq) + 4H^+(aq) + 2e^- \rightleftharpoons PbSO_4(s) + 2H_2O(l)$	+1.69
$MnO_4^-(aq) + 8H^+(aq) + 5e^- \rightleftharpoons Mn^{2+}(aq) + 4H_2O(l)$	+1.51
$Au^{3+}(aq) + 3e^- \rightleftharpoons Au(s)$	+1.50
$ClO_4^-(aq) + 8H^+(aq) + 8e^- \rightleftharpoons Cl^-(aq) + 4H_2O(l)$	+1.39
$Cl_2(g) + 2e^- \rightleftharpoons 2Cl^-(aq)$	+1.36
$2HNO_2(aq) + 4H^+(aq) + 4e^- \rightleftharpoons N_2O(g) + 3H_2O(l)$	+1.30
$Cr_2O_7^{2-}(aq) + 14H^+(aq) + 6e^- \rightleftharpoons 2Cr^{3+}(aq) + 7H_2O(l)$	+1.23
$O_2(g) + 4H^+(aq) + 4e^- \rightleftharpoons 2H_2O(l)$	+1.23
$MnO_2(s) + 4H^+(aq) + 2e^- \rightleftharpoons Mn^{2+}(aq) + 2H_2O(l)$	+1.22
$Br_2(l) + 2e^- \rightleftharpoons 2Br^-(aq)$	+1.07
$Hg^{2+}(aq) + 2e^- \rightleftharpoons Hg(l)$	+0.85
$OCl^-(aq) + H_2O(l) + 2e^- \rightleftharpoons Cl^-(aq) + 2OH^-(aq)$	+0.84
$2NO_3^-(aq) + 4H^+(aq) + 2e^- \rightleftharpoons N_2O_4(g) + 2H_2O(l)$	+0.80
$Ag^+(aq) + e^- \rightleftharpoons Ag(s)$	+0.80
$Fe^{3+}(aq) + e^- \rightleftharpoons Fe^{2+}(aq)$	+0.77
$O_2(g) + 2H^+(aq) + 2e^- \rightleftharpoons H_2O_2(l)$	+0.70
$I_2(s) + 2e^- \rightleftharpoons 2I^-(aq)$	+0.54
$O_2(g) + 2H_2O(l) + 4e^- \rightleftharpoons 4OH^-(aq)$	+0.40
$Cu^{2+}(aq) + 2e^- \rightleftharpoons Cu(s)$	+0.34
$SO_4^{2-}(aq) + 4H^+(aq) + 2e^- \rightleftharpoons H_2SO_3(aq) + H_2O(l)$	+0.17
$Sn^{4+}(aq) + 2e^- \rightleftharpoons Sn^{2+}(aq)$	+0.15
$S(s) + 2H^+(aq) + 2e^- \rightleftharpoons H_2S(aq)$	+0.14
$AgBr(s) + e^- \rightleftharpoons Ag(s) + Br^-(aq)$	+0.07
$2H^+(aq) + 2e^- \rightleftharpoons H_2(g)$	0.00
$Pb^{2+}(aq) + 2e^- \rightleftharpoons Pb(s)$	-0.13
$Sn^{2+}(aq) + 2e^- \rightleftharpoons Sn(s)$	-0.14
$AgI(s) + e^- \rightleftharpoons Ag(s) + I^-(aq)$	-0.15
$Ni^{2+}(aq) + 2e^- \rightleftharpoons Ni(s)$	-0.26
$Co^{2+}(aq) + 2e^- \rightleftharpoons Co(s)$	-0.28
$PbSO_4(s) + 2e^- \rightleftharpoons Pb(s) + SO_4^{2-}(aq)$	-0.36
$Se(s) + 2H^+(aq) + 2e^- \rightleftharpoons H_2Se(aq)$	-0.40
$Cd^{2+}(aq) + 2e^- \rightleftharpoons Cd(s)$	-0.40
$Cr^{3+}(aq) + e^- \rightleftharpoons Cr^{2+}(aq)$	-0.41
$Fe^{2+}(aq) + 2e^- \rightleftharpoons Fe(s)$	-0.45
$NO_2^-(aq) + H_2O(l) + e^- \rightleftharpoons NO(g) + 2OH^-(aq)$	-0.46
$Ag_2S(s) + 2e^- \rightleftharpoons 2Ag(s) + S^{2-}(aq)$	-0.69
$Zn^{2+}(aq) + 2e^- \rightleftharpoons Zn(s)$	-0.76
$2H_2O(l) + 2e^- \rightleftharpoons H_2(g) + 2OH^-(aq)$	-0.83
$Cr^{2+}(aq) + 2e^- \rightleftharpoons Cr(s)$	-0.91
$Se(s) + 2e^- \rightleftharpoons Se^{2-}(aq)$	-0.92
$SO_4^{2-}(aq) + H_2O(l) + 2e^- \rightleftharpoons SO_3^{2-}(aq) + 2OH^-(aq)$	-0.93
$Al^{3+}(aq) + 3e^- \rightleftharpoons Al(s)$	-1.66
$Mg^{2+}(aq) + 2e^- \rightleftharpoons Mg(s)$	-2.37
$Na^+(aq) + e^- \rightleftharpoons Na(s)$	-2.71
$Ca^{2+}(aq) + 2e^- \rightleftharpoons Ca(s)$	-2.87
$Ba^{2+}(aq) + 2e^- \rightleftharpoons Ba(s)$	-2.91
$K^+(aq) + e^- \rightleftharpoons K(s)$	-2.93
$Li^+(aq) + e^- \rightleftharpoons Li(s)$	-3.04

*For 1.0 mol/L solutions at 298.15 K (25.00 °C) and a pressure of 101.325 kPa

Relative Strengths of Acids and Bases at 298.15 K

Common Name IUPAC / Systematic Name	Acid Formula	Conjugate Base Formula	K_a
perchloric acid aqueous hydrogen perchlorate	$\text{HClO}_4(\text{aq})$	$\text{ClO}_4^-(\text{aq})$	very large
hydroiodic acid aqueous hydrogen iodide	$\text{HI}(\text{aq})$	$\text{I}^-(\text{aq})$	very large
hydrobromic acid aqueous hydrogen bromide	$\text{HBr}(\text{aq})$	$\text{Br}^-(\text{aq})$	very large
hydrochloric acid aqueous hydrogen chloride	$\text{HCl}(\text{aq})$	$\text{Cl}^-(\text{aq})$	very large
sulfuric acid aqueous hydrogen sulfate	$\text{H}_2\text{SO}_4(\text{aq})$	$\text{HSO}_4^-(\text{aq})$	very large
nitric acid aqueous hydrogen nitrate	$\text{HNO}_3(\text{aq})$	$\text{NO}_3^-(\text{aq})$	very large
hydronium ion	$\text{H}_3\text{O}^+(\text{aq})$	$\text{H}_2\text{O}(\text{l})$	1
oxalic acid	$\text{HOOC-COOH}(\text{aq})$	$\text{HOOC-COO}^-(\text{aq})$	5.6×10^{-2}
sulfurous acid aqueous hydrogen sulfite	$\text{H}_2\text{SO}_3(\text{aq})$	$\text{HSO}_3^-(\text{aq})$	1.4×10^{-2}
hydrogen sulfate ion	$\text{HSO}_4^-(\text{aq})$	$\text{SO}_4^{2-}(\text{aq})$	1.0×10^{-2}
phosphoric acid aqueous hydrogen phosphate	$\text{H}_3\text{PO}_4(\text{aq})$	$\text{H}_2\text{PO}_4^-(\text{aq})$	6.9×10^{-3}
citric acid 2-hydroxy-1,2,3-propanetricarboxylic acid	$\text{C}_3\text{H}_5\text{O}(\text{COOH})_3(\text{aq})$	$\text{C}_3\text{H}_5\text{O}(\text{COOH})_2\text{COO}^-(\text{aq})$	7.4×10^{-4}
hydrofluoric acid aqueous hydrogen fluoride	$\text{HF}(\text{aq})$	$\text{F}^-(\text{aq})$	6.3×10^{-4}
nitrous acid aqueous hydrogen nitrite	$\text{HNO}_2(\text{aq})$	$\text{NO}_2^-(\text{aq})$	5.6×10^{-4}
formic acid methanoic acid	$\text{HCOOH}(\text{aq})$	$\text{HCOO}^-(\text{aq})$	1.8×10^{-4}
hydrogen oxalate ion	$\text{HOOC-COO}^-(\text{aq})$	$\text{OOC-COO}^{2-}(\text{aq})$	1.5×10^{-4}
lactic acid 2-hydroxypropanoic acid	$\text{C}_2\text{H}_5\text{O-COOH}(\text{aq})$	$\text{C}_2\text{H}_5\text{O-COO}^-(\text{aq})$	1.4×10^{-4}
ascorbic acid 2(1,2-dihydroxyethyl)-4,5-dihydroxy-furan-3-one	$\text{H}_2\text{C}_6\text{H}_6\text{O}_6(\text{aq})$	$\text{HC}_6\text{H}_6\text{O}_6^-(\text{aq})$	9.1×10^{-5}

benzoic acid benzenecarboxylic acid	$C_6H_5COOH(aq)$	$C_6H_5COO^-(aq)$	6.3×10^{-5}
acetic acid ethanoic acid	$CH_3COOH(aq)$	$CH_3COO^-(aq)$	1.8×10^{-5}
dihydrogen citrate ion	$C_3H_5O(COOH)_2COO^-(aq)$	$C_3H_5O(COOH)(COO)_2^{2-}(aq)$	1.7×10^{-5}
butanoic acid	$C_3H_7COOH(aq)$	$C_3H_7COO^-(aq)$	1.5×10^{-5}
propanoic acid	$C_2H_5COOH(aq)$	$C_2H_5COO^-(aq)$	1.3×10^{-5}
carbonic acid ($CO_2 + H_2O$) aqueous hydrogen carbonate	$H_2CO_3(aq)$	$HCO_3^-(aq)$	4.5×10^{-7}
hydrogen citrate ion	$C_3H_5O(COOH)(COO)_2^{2-}(aq)$	$C_3H_5O(COO)_3^{3-}(aq)$	4.0×10^{-7}
hydrosulfuric acid aqueous hydrogen sulfide	$H_2S(aq)$	$HS^-(aq)$	8.9×10^{-8}
hydrogen sulfite ion	$HSO_3^-(aq)$	$SO_3^{2-}(aq)$	6.3×10^{-8}
dihydrogen phosphate ion	$H_2PO_4^-(aq)$	$HPO_4^{2-}(aq)$	6.2×10^{-8}
hypochlorous acid aqueous hydrogen hypochlorite	$HOCl(aq)$	$OCl^-(aq)$	4.0×10^{-8}
hydrocyanic acid aqueous hydrogen cyanide	$HCN(aq)$	$CN^-(aq)$	6.2×10^{-10}
ammonium ion	$NH_4^+(aq)$	$NH_3(aq)$	5.6×10^{-10}
hydrogen carbonate ion	$HCO_3^-(aq)$	$CO_3^{2-}(aq)$	4.7×10^{-11}
hydrogen ascorbate ion	$HC_6H_6O_6^-(aq)$	$C_6H_6O_6^{2-}(aq)$	2.0×10^{-12}
hydrogen phosphate ion	$HPO_4^{2-}(aq)$	$PO_4^{3-}(aq)$	4.8×10^{-13}
water	$H_2O(l)$	$OH^-(aq)$	1.0×10^{-14}

Note: An approximation may be used instead of the quadratic formula when the concentration of H_3O^+ produced is less than 5% of the original acid concentration (or the concentration of the acid is 1 000 times greater than the K_a). An approximation can also be used for weak bases. The formulas of the carboxylic acids have been written so that the COOH group can be easily recognized. Either the common or IUPAC name is acceptable.

Acid–Base Indicators at 298.15 K

Indicator	Suggested Abbreviations	pH Range	Colour Change as pH Increases	K_a
methyl violet	HMv(aq) / Mv ⁻ (aq)	0.0 – 1.6	yellow to blue	$\sim 2 \times 10^{-1}$
cresol red	H ₂ Cr(aq) / HCr ⁻ (aq)	0.0 – 1.0	red to yellow	$\sim 3 \times 10^{-1}$
	HCr ⁻ (aq) / Cr ²⁻ (aq)	7.0 – 8.8	yellow to red	3.5×10^{-9}
thymol blue	H ₂ Tb(aq) / HTb ⁻ (aq)	1.2 – 2.8	red to yellow	2.2×10^{-2}
	HTb ⁻ (aq) / Tb ²⁻ (aq)	8.0 – 9.6	yellow to blue	6.3×10^{-10}
orange IV	HOr(aq) / Or ⁻ (aq)	1.4 – 2.8	red to yellow	$\sim 1 \times 10^{-2}$
methyl orange	HMo(aq) / Mo ⁻ (aq)	3.2 – 4.4	red to yellow	3.5×10^{-4}
bromocresol green	HBg(aq) / Bg ⁻ (aq)	3.8 – 5.4	yellow to blue	1.3×10^{-5}
methyl red	HMr(aq) / Mr ⁻ (aq)	4.8 – 6.0	red to yellow	1.0×10^{-5}
chlorophenol red	HCh(aq) / Ch ⁻ (aq)	5.2 – 6.8	yellow to red	5.6×10^{-7}
bromothymol blue	HBb(aq) / Bb ⁻ (aq)	6.0 – 7.6	yellow to blue	5.0×10^{-8}
phenol red	HPr(aq) / Pr ⁻ (aq)	6.6 – 8.0	yellow to red	1.0×10^{-8}
phenolphthalein	HPh(aq) / Ph ⁻ (aq)	8.2 – 10.0	colourless to pink	3.2×10^{-10}
thymolphthalein	HTh(aq) / Th ⁻ (aq)	9.4 – 10.6	colourless to blue	1.0×10^{-10}
alizarin yellow R	HAy(aq) / Ay ⁻ (aq)	10.1 – 12.0	yellow to red	6.9×10^{-12}
indigo carmine	HIc(aq) / Ic ⁻ (aq)	11.4 – 13.0	blue to yellow	$\sim 6 \times 10^{-12}$
1,3,5–trinitrobenzene	HNb(aq) / Nb ⁻ (aq)	12.0 – 14.0	colourless to orange	$\sim 1 \times 10^{-13}$

Colours of Common Aqueous Ions

Ionic Species	Solution Concentration	
	1.0 mol/L	0.010 mol/L
chromate	yellow	pale yellow
chromium(III)	blue-green	green
chromium(II)	dark blue	pale blue
cobalt(II)	red	pink
copper(I)	blue-green	pale blue-green
copper(II)	blue	pale blue
dichromate	orange	pale orange
iron(II)	lime green	colourless
iron(III)	orange-yellow	pale yellow
manganese(II)	pale pink	colourless
nickel(II)	blue-green	pale blue-green
permanganate	deep purple	purple-pink

